

Solution set 4

to some problems given for TMA4230 Functional analysis

2004-03-17

Exercise A.1. I will not write up the full solutions to all of these simple exercises.

To prove $\bigcup\{C^c: C \in \mathcal{C}\} = (\bigcap\mathcal{C})^c$, note that $x \in \bigcup\{C^c: C \in \mathcal{C}\}$ means $x \in C^c$ for every $C \in \mathcal{C}$, which means $x \notin C$ for every $C \in \mathcal{C}$, which means $x \in C$ for no $C \in \mathcal{C}$, which is the opposite of $x \in C$ for some $C \in \mathcal{C}$, which means $x \notin \bigcap\mathcal{C}$.

And to prove $f^{-1}(A \setminus B) = f^{-1}(A) \setminus f^{-1}(B)$, note that $x \in f^{-1}(A \setminus B)$ means $f(x) \in A \setminus B$, which means $f(x) \in A$ but $f(x) \notin B$, which is the same as $x \in f^{-1}(A)$ but $x \notin f^{-1}(B)$, which is the same as $x \in f^{-1}(A) \setminus f^{-1}(B)$.

Exercise A.2. Remember that f is continuous if and only if $f^{-1}(V)$ is open for every open $V \subseteq Y$, and note that taking complements establishes a one-to-one correspondence between open sets and closed sets. So f is continuous if and only if $f^{-1}(V)^c$ is closed for every open $V \subseteq Y$, which is equivalent to $f^{-1}(V^c)$ being closed for every open $V \subseteq Y$, which is equivalent to $f^{-1}(F)$ being closed for every closed $F \subseteq Y$ (where we replaced V by F^c in the last step).

Exercise A.3. First, assume that K is compact with the original definition. Let \mathcal{V} be an open cover of K in the sense of the last paragraph of the exercise. Then let $\mathcal{U} = \{K \cap V: V \in \mathcal{V}\}$. Each member of \mathcal{U} is a (relatively) open subset of K , and $\bigcup\mathcal{U} = K \cap \bigcup V = K$, since $\bigcup V \supseteq K$. Thus \mathcal{U} is an open cover of K in the original definition. By compactness of K , it has a finite subcover $\{K \cap V_1, \dots, K \cap V_n$ with $V_1, \dots, V_n \in \mathcal{V}$. So $(K \cap V_1) \cup \dots \cup (K \cap V_n) = K$, which implies $\{V_1 \cup \dots \cup V_n \supset K$, and one implication is proved.

Conversely, if K satisfies the condition of the last paragraph of the exercise, let \mathcal{U} be an open cover of K as originally defined. Let \mathcal{V} consist of all open subsets $V \subseteq X$ so that $K \cap V \in \mathcal{U}$. Since every $U \in \mathcal{U}$ is open, it can be written $K \cap V$ for at least one open set $V \subseteq X$, and so \mathcal{V} is an open cover (new definition) for K . Pick a finite subcover (new definition) $\{V_1, \dots, V_n\}$. Then $\{K \cap V_1, \dots, K \cap V_n\}$ is a finite subcover (original definition) of K .

Exercise A.4. Let K be compact, and let $L \subseteq K$ be closed. Any collection of closed subsets of L is also a collection of closed subsets of K , and if it has the finite intersection property, it has nonempty intersection because K is compact. This shows that L is compact.

Exercise A.5. Let X be Hausdorff and K a compact subset of X .

If $x \in \overline{K}$, there is a filter \mathcal{F} in K converging to x . (In fact, one can take \mathcal{F} to be generated by the sets $K \cap V$ for neighbourhoods V of x .) Since K is compact, there is a finer filter \mathcal{G} which converges in K , to a point $y \in K$. This filter then converges to y in X as well, and it converges to x too, since it refines \mathcal{F} . But no filter can have two distinct limits on a Hausdorff space, so $x = y \in K$. Thus K is closed. (There are many different ways to organize this proof, not all of which use filters.)

Exercise A.6.

(a) First, if $x \neq y$ then the Hausdorff property means that x and y have disjoint neighbourhoods in the original topology. But then these are neighbourhoods in the stronger topology as well, and the Hausdorff property in the stronger topology follows.

Now let $F \subseteq X$ be closed in the stronger topology, but not in the original topology. If X is compact in the stronger topology, then so is F (exercise A.4). But then F is compact in the original topology as well, by the easy part of (b) below. This implies that F is closed in the original topology (exercise A.5), and this is a contradiction.

(b) First, consider an open (in the weaker topology) cover of X . Then this is an open (in the original topology) cover as well, and by compactness in the original topology, it has a finite subcover. So X is compact in the weaker topology.

Now let $F \subseteq X$ be closed in the original topology, but not in the weaker topology. Consider an open cover of F (in the weaker topology, and in the sense of exercise A.3). This is then also an open cover of F in the original topology, so it has a finite subcover. This shows that F is in fact compact in the weaker topology. If the weaker topology is Hausdorff, then F must be closed in the weaker topology (exercise A.5), and this is a contradiction.

Exercise A.7. By definition, the constructed collection \mathcal{T} contains X as a member, and arbitrary unions of elements of \mathcal{T} belong to \mathcal{T} . To show that \mathcal{T} is a topology, it only remains to show that $U \cap V \in \mathcal{T}$ when $U, V \in \mathcal{T}$. So write $U = \bigcup \mathcal{U}$ where $\mathcal{U} \subseteq \mathcal{B}'$, and $V = \bigcup \mathcal{V}$ where $\mathcal{V} \subseteq \mathcal{B}'$. Then

$$U \cap V = (\bigcup \mathcal{U}) \cap (\bigcup \mathcal{V}) = \bigcup_{U \in \mathcal{U}} (U \cap \bigcup \mathcal{V}) = \bigcup_{U \in \mathcal{U}} \bigcup_{V \in \mathcal{V}} U \cap V \in \mathcal{T},$$

because each U and each V in the union belongs to \mathcal{B}' , and so $U \cap V \in \mathcal{B}'$ as well.

Because \mathcal{T} is created from \mathcal{B} by taking finite intersections and arbitrary unions, it is clear that any topology containing \mathcal{B} must contain \mathcal{T} as well. Thus \mathcal{T} is the weakest topology containing \mathcal{B} , as stated.

My definitions of *base* for a topology was wrong. (Even the word was wrong: It should be *base*, not *basis*.) A base \mathcal{B} for a topology \mathcal{T} is a subset of \mathcal{T} so that every member of \mathcal{T} is a union of members of \mathcal{B} . In the above situation, \mathcal{B}' is a base for \mathcal{T} , but \mathcal{B} need not be. This has no consequences for the next problem, for \mathcal{B}' is countable if \mathcal{B} is countable.

Exercise A.8. In a metric space X , any point x has the countable filterbase consisting of balls $B_{1/n}(x)$, for $n = 1, 2, \dots$

If \mathcal{T} has a countable base \mathcal{B} and $x \in X$, then $\{V \in \mathcal{B} : x \in V\}$ is a countable base for the neighbourhood filter at x , so X is first countable.

It is tempting to look for a metrizable space that is not second countable, since metric spaces are automatically first countable. In fact we may try using the discrete topology on some set X , which is first countable in the extreme ($\{\{x\}\}$ is a base for the neighbourhood filter at x). If X is uncountable, then surely the discrete topology on X cannot be second countable?

This seems to require a bit more set theory than I had realized when I posed this question, so I am afraid I have been a bit unfair.

Let us go a bit further and let $X = \mathbb{R}$. Remember, we are using the *discrete* topology on X – I am choosing the set \mathbb{R} only because this set has lots of elements. A topology \mathcal{T} with a countable basis \mathcal{B} has a cardinality at most equal to that of \mathbb{R} , since it is formed by unions of members of the countable set \mathcal{B} , and the set of subsets of a countably infinite sets has cardinality equal to that of \mathbb{R} . But the discrete topology on X is the set of *all* subsets of X , and that has greater cardinality than X itself, and hence greater cardinality to \mathcal{T} . So the discrete topology on \mathbb{R} is not second countable.

What is cardinality? Recall that any set X can be wellordered. With each such wellorder, X becomes order isomorphic to some ordinal number. The cardinality of X is the smallest ordinal number one can get in this way. A *cardinal number* is an ordinal number which is the cardinality of some set.

For example, every finite ordinal number is a cardinal number, and so is ω , the smallest infinite ordinal. ω is the cardinality of the natural numbers. When we think of ω as a cardinal number, we also write it as \aleph_0 .¹ The immediate successors $\omega + 1$, $\omega + 2$, \dots , $\omega + \omega$ and so forth are not cardinal numbers, since they are all countable. The smallest cardinal number bigger than \aleph_0 is called \aleph_1 , and so forth and so on.

Cantor's theorem states that the set $\mathcal{P}(X)$ of subsets of a given set X has strictly greater cardinality than X itself. In particular the cardinality of $\mathcal{P}(\mathbb{N})$ is greater than \aleph_0 . What used to be a famous conjecture is that the cardinality of $\mathcal{P}(\mathbb{N})$, also written as 2^{\aleph_0} , equals \aleph_1 . This so-called *continuum hypothesis* has later been shown to be independent of the other axioms of set theory, as has the *generalized continuum hypothesis*, that $2^{\aleph_n} = \aleph_{n+1}$ for every ordinal n .

The proof of Cantor's theorem is by the famous *diagonal argument*: Assume that $f: X \rightarrow \mathcal{P}(X)$ maps X onto $\mathcal{P}(X)$. Form the set $D = \{x \in X : x \notin f(x)\}$. Now if $D = f(w)$ for some $w \in X$, then $w \in D \Leftrightarrow w \notin D$, a contradiction.

¹ \aleph is the first letter of the Hebrew alphabet. A vowel, in a written language without vowels? I am not sure I understand this.